

Lattice Exploration of Strong Dynamics at the TeV Scale

Meifeng Lin



Brookhaven Forum 2011
October 19 - 21, 2011

Outline

Motivation

Lattice Calculations

Lattice search for the conformal window
Work by the LSD Collaboration

Summary

Technicolor Theories

Weinberg & Susskind 1979

- ▶ **Strong Dynamics:**
strongly coupled gauge theories *may* play a role at the electroweak scale: $\Lambda_{TC} \simeq F_{EW} \approx 246 \text{ GeV}$.
- ▶ Introduce new gauge interactions $SU(N_{TC})$ at Λ_{TC} .
- ▶ Technifermions possess chiral symmetry, transforming under $SU(N_{f,TC})_L \otimes SU(N_{f,TC})_R$.
- ▶ Spontaneous chiral symmetry breaking
→ massless Technicolor Goldstone bosons
→ three of these provide mass for the EW gauge bosons.
- ▶ Extended technicolor at $\Lambda_{ETC} \gg \Lambda_{TC}$ to generate quark and lepton masses. Dimopoulos & Susskind 1979, Eichten & Lane 1979

$$m_{q,l} \simeq \frac{\langle \bar{Q}Q \rangle_{ETC}}{\Lambda_{ETC}^2}$$

Difficulties with QCD-like Technicolor

If TC theories are QCD-like

- ▶ With $\langle \bar{Q}Q \rangle_{TC} \simeq 4\pi F_{EW}^3$, and

$$\langle \bar{Q}Q \rangle_{ETC} = \langle \bar{Q}Q \rangle_{TC} \exp \left(\int_{\Lambda_{TC}}^{\Lambda_{ETC}} \frac{d\mu}{\mu} \gamma_m(\mu) \right) \rightarrow \langle \bar{Q}Q \rangle_{TC} \ln(\Lambda_{ETC}/\Lambda_{TC})$$

$$\rightarrow \langle \bar{Q}Q \rangle_{ETC} / \Lambda_{ETC}^2 < 0.1 \text{ MeV}$$

→ too small to generate SM quark masses

- ▶ The electroweak S parameter would be too big to be consistent with the experimental constraint: $S \approx 0$.

One solution: **Walking Technicolor**.

Walking Technicolor

There exists a region $\Lambda_{IR} < \mu < \Lambda_{UV}$ where the running coupling $\alpha(\mu)$ evolves very slowly.

- ▶ Assume $\gamma(\mu) \sim \gamma$, then Techni-quark condensate gets enhanced:

$$\langle \bar{Q}Q \rangle_{ETC} \sim \langle \bar{Q}Q \rangle_{TC} \left(\frac{\Lambda_{UV}}{\Lambda_{IR}} \right)^\gamma$$

Could be large enough to generate quark masses.

- ▶ Could result in a small S parameter.

Such theories could exist with a large number of massless fermions.

The Conformal Window

- ▶ SU(N) gauge theories with N_f massless Dirac fermions, conjectured from two-loop β function

Banks & Zaks 1981

$N_f < 11N/2$	asymptotically free
$0 \leq N_f < N_f^c$	confinement and S χ SB
$N_f^c \leq N_f < 11N/2$	conformal (\exists Infrared fixed point)
$N_f > 11N_c/2$	asymptotic freedom lost

- ▶ N_f^c – the conformal window
- ▶ “Walking” may appear with $N_f \rightarrow N_f^c$.
- ▶ The problem: N_f^c not precisely known. Various perturbative calculations and approximations give a wide range of results.
- ▶ Lattice calculations can help.

Lattice Toolbox

Many existing tools for the QCD studies can be used:

- ▶ Non-perturbative running coupling calculations
How does the running coupling change with the energy scale?
- ▶ Hadron and glueball spectrum measurements
Are the quark mass dependence and state ordering the same as QCD?
- ▶ Chiral condensate
Does the chiral condensate vanish or get enhanced?
- ▶ Thermodynamic properties (T_c , EoS, etc.)
Is there a deconfinement transition? Is EoS different?

A Dozen Lattice BSM Efforts Worldwide



Aoyama et al.



DeGrand et al.



Del Debbio et al.



Deuzeman et al.



Catteral et al.



LSD



Hietanen et al.



A. Hasenfratz



LHC



Jin-Mawhinney




Yamada et al.




Kogut-Sinclair

Picture Credit: George Fleming (Origin of Mass 2010)

Current Landscape



(Updated 07/12/11)



Picture Credit: Ethan Neil (Lattice 2011)

The Lattice Strong Dynamics Collaboration

20 people from 9 institutions:

Tom Appelquist Yale U.

Ron Babich Boston U.

Rich Brower Boston U.

Michael Buchoff LLNL

Michael Cheng Boston U.

Mike Clark Harvard U.

Saul Cohen U. Washington

George Fleming Yale U.

Joe Kiskis U. of California, Davis

Meifeng Lin Yale U.

Heechang Na ANL

Ethan Neil Fermilab

James Osborn ANL

Claudio Rebbi Boston U.

David Schaich Boston U.

Sergey Syritsyn LBL

Joe Wasem LLNL

Oliver Witzel Boston U.

Gennady Voronov Yale U.

Pavlos Vranas LLNL

LSD Program

- ▶ Explore the $N_c - N_f$ plane of the $SU(N_c)$ gauge theories non-perturbatively.
- ▶ Look for trends with changing N_c , N_f , and **perhaps** locate N_f^c .
- ▶ Chiral symmetry plays an important role.
 - ▶ Use domain wall fermions: nearly exact chiral symmetry.
- ▶ Start from something familiar on the lattice: $SU(3)$ in the fundamental representation.
 - ▶ $N_f = 2$: as a reference point.
 - ▶ $N_f = 6$: first focus on QCD-like theories, and move cautiously towards the conformal window.
 - ▶ $N_f = 10$: more challenging, can be QCD-like, conformal or walking.
- ▶ Want to have sufficient separation between IR and UV scales in a slowly-running theory \rightarrow small lattice spacing:
 - ▶ **We aim for five times the vector meson mass: $1/a \approx 5M_V$.**

Current Status

- ▶ $N_f = 2$ and $N_f = 6$, more than three years of running. Currently increasing statistics on existing ensembles.
- ▶ Observables under study:
 - ▶ hadron spectrum and chiral condensate
 - ▶ S parameter
 - ▶ $\pi - \pi$ scattering
- ▶ Publications for $N_f = 2$ and $N_f = 6$:
 - ▶ Condensate enhancement, arXiv:0910.2224 [hep-ph].
 - ▶ S parameter and parity doubling, arXiv:1009.5967 [hep-ph]
- ▶ $N_f = 10$ in progress.
- ▶ $SU(2)$ simulations in the pipeline.

Chiral Condensate Enhancement

- ▶ As we increase N_f but below N_f^c , if the theory becomes slowly running, then the chiral condensate should get enhanced.
- ▶ On the lattice we can use three ways to measure chiral condensate (from three GMOR ratios)


$$R = \frac{\langle \bar{\psi}\psi \rangle}{F_\pi^3} = \frac{M_\pi^3}{\sqrt{(2m)^3 \langle \bar{\psi}\psi \rangle}} = \frac{M_\pi^2}{2mF_\pi}$$

- ▶ Define enhancement ratios w.r.t the SM with $N_f = 2$ **at roughly the same lattice cutoff**.

$$\mathcal{R}_{XY, \tilde{m}} \equiv \frac{R^{(N_f)}}{R^{(N_f=2)}}$$

Condensate Enhancement for $N_f = 6$

$$R_m \equiv \frac{M_m^2}{2mF_\pi(N_f=6)} / \frac{M_m^2}{2mF_\pi(N_f=2)}$$



T. Appelquist *et al.* (LSD Collaboration), PRL104:071601,2010

The S Parameter

- ▶ Parametrizes vacuum polarization (oblique) corrections. Peskin and Takeuchi 1992
- ▶ Used to constrain new physics beyond the Standard Model.
- ▶ Definition

$$S = -4\pi [\Pi'_{VV}(0) - \Pi'_{AA}(0)] - \Delta S_{SM}, \quad \Pi'(0) = \frac{d\Pi(q^2)}{dq^2} \Big|_{q^2 \rightarrow 0}$$

where

$$\Pi_{VV}^{\mu\nu}(q) = \sum_x e^{iq \cdot x} \langle V^\mu(x) V^\nu(0) \rangle, \quad \Pi_{AA}^{\mu\nu}(q) = \sum_x e^{iq \cdot x} \langle A^\mu(x) A^\nu(0) \rangle$$


- ▶ ΔS_{SM} – Standard Model Higgs contributions

$$\Delta S_{SM} = \frac{1}{4} \int_0^\infty \frac{ds}{s} \left[1 - \left(1 - \frac{m_H^2}{s} \right)^3 \theta(s - m_H^2) \right]$$

m_H – reference Higgs mass.

- ▶ Electroweak precision experiments find $S \approx 0$.
Scaled-up QCD with $N_f = 2$ gives $S \approx 0.3$.

Extracting S from Current Correlators



Legend:

◆ $N_f=2$

▲ $N_f=6$

■ $m=0.005$

■ $m=0.010$

■ $m=0.015$

■ $m=0.020$


■ $m=0.025$

■ $m=0.030$


- ▶ Need the slopes at $q^2 = 0$.
- ▶ Use Padè(1,2) to describe the q^2 dependence.

$$\Pi_{V-A}(q^2) = \frac{a_0 + a_1 q^2}{1 + q^2(b_1 + b_2 q^2)}$$

- ▶ Fits describe the data reasonably well in the small q^2 region
- ▶ Insensitive to fit ranges: fits are stable.



The S Parameter



T. Appelquist *et al.* (LSD Collaboration), PRL 106 (2011) 231601

- ▶ $N_f = 2$ gives $S = 0.32(5)$ with $m_H \sim 1$ TeV, consistent with scaled-up QCD.
- ▶ For $N_f = 6$, at small mass, S drops below naive scaled-up QCD value, which would be 3 times the $N_f = 2$ value.
- ▶ There can be chiral log contributions, which will eventually make S turn up. (In $N_f = 2$, the chiral logs cancel after the SM subtraction.)
- ▶ Other interactions need to be turned on to have a finite S for $N_f = 6$ in the chiral limit.

Is the ten-flavor theory chirally broken, or conformal?

- ▶ Hayakawa *et al.* observed evidence for the existence of an infrared fixed point from Schrödinger functional running coupling studies. [Phys. Rev. D 83, 074509 (2011)]
- ▶ The hadron mass quark mass dependence is different in these two scenarios.
- ▶ In a **chirally broken** theory, at sufficiently light quark masses, chiral perturbation theory predicts m_f dependence.
 - ▶ However, M_p/M_V values in our simulations range from 0.71 to 0.83.
 - ▶ Simulated quark masses are likely too heavy to make use of ChPT.
- ▶ In a **mass-deformed conformal** theory, hadron masses are governed by universality,

$$M_X \approx C_X m^{1/(1+\gamma^*)} + \dots$$


- ▶ Ratios of hadron masses would be roughly independent of quark masses.
- ▶ For DWF, $m \equiv m_f + m_{\text{res}}$.

Can the theory be conformal?

Recall that in a mass-deformed conformal theory

$$M_X = C_X m^{1/(1+\gamma^*)}$$

so the vector-to-pseudoscalar meson mass ratio M_V/M_P should look roughly constant.



- ▶ The $N_f = 2$ data are diverging as $m \rightarrow 0$, as expected since $M_P \rightarrow 0$.
- ▶ Similarly for $N_f = 6$, though not as obvious.
- ▶ The $N_f = 10$ data may show an upward trend as $m \rightarrow 0$. But since the errorbars are underestimated, it is also possible that the ratios can be roughly constant. **More work is in progress.**

Summary

- ▶ Lattice studies by the LSD collaboration find evidence for condensate enhancement and reduced S parameter with $N_f = 6$ for SU(3) gauge theories.
- ▶ $N_f = 10$ studies are in progress. Could be in either the chirally broken or conformal phase.
- ▶ Lattice gauge theory has the potential to provide valuable input to theoretical model building.

Challenges

- ▶ Computationally more demanding than QCD.
- ▶ There are still many unresolved issues that need to be understood.
- ▶ What is the best way to distinguish conformal from QCD-like?
- ▶ How do we distinguish walking from conformal? Larger boxes and smaller lattice spacings?